

**GLP-1 Receptor Agonist Treatment of Diabetes - Agreement Form and Checklist**  
*Use alongside [GLP-1 Receptor Agonist Guidelines](#). For Semaglutide, liraglutide and dulaglutide ONLY.*

<b>Date seen:</b>	<b>DSN name:</b>	
<b>Name:</b>	<b>GP name:</b>	
<b>NHS number:</b>	<b>GP Practice:</b>	
<b>DOB:</b>		
<b>Exclusion criteria:</b>	<div>Yes</div> <div>No</div>	<b>Warnings:</b>
<b>If YES do not prescribe or see note in red</b>		<div>YES</div> <div>NO</div>
eGFR <15ml/min (within last 2 mths) _____ml/min <b>AVOID ALL if &lt; 15ml/min</b>	<input type="checkbox"/> <input type="checkbox"/>	Person able to inject or be injected? If not, consider oral semaglutide. <input type="checkbox"/> <input type="checkbox"/>
Triglycerides >5MMOL/L in last 2 months _____mmol/l <b>CONSIDER RISK IF ABOVE 5MMOL/L</b>	<input type="checkbox"/> <input type="checkbox"/>	Person warned of possible side effects (e.g. GI, hypos, DKA)? Oral semaglutide must be taken on an empty stomach 30 mins before food. <input type="checkbox"/> <input type="checkbox"/>
Alcohol intake above recommended _____units/wk <b>If above national recommendations discuss increased pancreatitis risk</b>	<input type="checkbox"/> <input type="checkbox"/>	DKA <a href="#">MHRA warning</a> discussed <input type="checkbox"/> <input type="checkbox"/>
Biliary or pancreatic surgery or disease within last 6 months	<input type="checkbox"/> <input type="checkbox"/>	Person warned of risk of dehydration? <input type="checkbox"/> <input type="checkbox"/>
Any history of pancreatitis	<input type="checkbox"/> <input type="checkbox"/>	Person warned of risk of pancreatitis? <input type="checkbox"/> <input type="checkbox"/>
Heart Failure NYHA class IV	<input type="checkbox"/> <input type="checkbox"/>	Instructions given to seek prompt advice if experiencing persistent severe abdominal pain with or without vomiting? <input type="checkbox"/> <input type="checkbox"/>
Inflammatory bowel disease	<input type="checkbox"/> <input type="checkbox"/>	Sick day rules discussed <input type="checkbox"/> <input type="checkbox"/>
Gastroparesis	<input type="checkbox"/> <input type="checkbox"/>	Contraception discussed <input type="checkbox"/> <input type="checkbox"/>
Planned pregnancy, inadequate contraception or breastfeeding	<input type="checkbox"/> <input type="checkbox"/>	HCP aware of 1) risk of DKA if insulin dose <u>reduced too quickly</u> when GLP1RA started 2) risk of hypos if insulin/sulfonylurea <u>dose not reduced enough</u> when GLP1RA started
<b>Caution in semaglutide only</b>		<b>GLP-1 chosen:</b>
Retinopathy grade R2/R3/M1/P1	<input type="checkbox"/> <input type="checkbox"/>	<ul style="list-style-type: none"> <li>▪ Semaglutide (subcut)</li> <li>▪ Dulaglutide</li> <li>▪ Liraglutide</li> <li>▪ Semaglutide (oral)</li> </ul>
<b>If YES discuss with specialist team</b>		

<b>Advised re NICE 6 month review criteria:</b>	<input type="checkbox"/>	<input type="checkbox"/>
<b>To take away:</b> Written information	<input type="checkbox"/>	<input type="checkbox"/>
<b>DVLA information:</b>	<input type="checkbox"/>	<input type="checkbox"/>

Liraglutide/Semaglutide/Dulaglutide has been prescribed for you to help reduce your blood glucose levels and help you to lose weight.

Liraglutide/Semaglutide/Dulaglutide is not of benefit to everyone and the National Institute for Health and Care Excellence (NICE) have advised that treatment should only be continued for people who have a reasonable benefit; defined by NICE as a reduction in HbA1c (long term glucose reading) of 1% point (DCCT units) or 11mmol/mol (IFCC units) or more and a reduction in weight of 3% or more after 6 months of treatment.

Over the next 6 months we will monitor your HbA1c and weight to assess if you are one of the people who benefit from the treatment. Liraglutide/Semaglutide/Dulaglutide will only usually be continued beyond 6 months in people who have a reasonable benefit from the treatment as stated above.

In occasional cases, where there has been a large reduction in either HbA1c or weight (above that suggested by NICE) but not in both, we may still decide that it is beneficial to continue using Liraglutide/Semaglutide/Dulaglutide.

Prior to commencing therapy:		At 3 months:		At 6 months:		
Date:		Date:		Date:		
HbA1c						
Wt (kg)						
BMI (kg/m2)						
eGFR (ml/min/1.73m2)						
%age weight loss 3 & 6 months				Target weight loss of at least 3% achieved?	YES	NO
HbA1c reduction 3 & 6 months				Target HbA1c reduction of at least 11mmol/mol (IFCC units) or 1% point (DCCT units) achieved?	YES	NO

**Agreement**

The information above has been explained to me and I understand that treatment with Liraglutide/SC Semaglutide/Dulaglutide/oral Semaglutide should be discontinued after 6 months if the medicine does not appear to be having a reasonable benefit.

Name .....

Signature .....